

AD-A046 630

GEORGE WASHINGTON UNIV WASHINGTON D C INST FOR MANAG--ETC F/G 12/1  
ON SOME GOODNESS OF FIT TESTS FOR THE WEIBULL DISTRIBUTION WITH--ETC(U)  
OCT 77 M CHANDRA, N D SINGPURWALLA N00014-77-C-0263

UNCLASSIFIED


SERIAL-T-363

NL

| OF |  
ADA  
046630



END  
DATE  
FILMED  
12-77  
DDC



NATIONAL BUREAU OF STANDARDS  
MICROCOPY RESOLUTION TEST CHART

AD A 0 46630

12  
B.S.

THE  
GEORGE  
WASHINGTON  
UNIVERSITY

STUDENTS FACULTY STUDY R  
ESEARCH DEVELOPMENT FUT  
URE CAREER CREATIVITY CC  
MMUNITY LEADERSHIP TECH  
NOLOGY FRONTIER DESIGN  
ENGINEERING APP ENC  
GEORGE WASHINGTON UNIV



DDC  
RECEIVED  
NOV 21 1977  
F97

AD No. \_\_\_\_\_  
DDC FILE COPY

INSTITUTE FOR MANAGEMENT  
SCIENCE AND ENGINEERING  
SCHOOL OF ENGINEERING  
AND APPLIED SCIENCE

THIS DOCUMENT HAS BEEN APPROVED FOR PUBLIC RELEASE AND SALE; ITS DISTRIBUTION IS UNLIMITED

6 ON SOME GOODNESS OF FIT TESTS FOR THE WEIBULL  
DISTRIBUTION WITH ESTIMATED PARAMETERS .

12  
by


10 Mahesh/Chandra  
Nozer D./Singpurwalla

9 Scientific rept.,

14 Serial-T-363  
14 October 1977

11 14 Oct 77

12 22p.



The George Washington University  
School of Engineering and Applied Science  
Institute for Management Science and Engineering

15  
Contract N00014-77-C-0263  
Project NR 042-372  
Office of Naval Research

This document has been approved for public  
sale and release; its distribution is unlimited.

406 743

1B

NONE

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER T-363	2. GOVT ACCESSION NO.	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle) ON SOME GOODNESS OF FIT TESTS FOR THE WEIBULL DISTRIBUTION WITH ESTIMATED PARAMETERS		5. TYPE OF REPORT & PERIOD COVERED SCIENTIFIC
7. AUTHOR(s) MAHESH CHANDRA NOZER D. SINGPURWALLA		6. PERFORMING ORG. REPORT NUMBER
9. PERFORMING ORGANIZATION NAME AND ADDRESS THE GEORGE WASHINGTON UNIVERSITY INSTITUTE FOR MANAGEMENT SCIENCE AND ENGINEERING WASHINGTON, D. C. 20037		8. CONTRACT OR GRANT NUMBER(s) N00014-77-C-0263 <i>New</i>
11. CONTROLLING OFFICE NAME AND ADDRESS OFFICE OF NAVAL RESEARCH CODE 436 ARLINGTON, VIRGINIA 22217		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)		12. REPORT DATE 14 October 1977
		13. NUMBER OF PAGES 18
		15. SECURITY CLASS. (of this report) NONE
16. DISTRIBUTION STATEMENT (of this Report) DISTRIBUTION OF THIS DOCUMENT IS UNLIMITED.		15a. DECLASSIFICATION/DOWNGRADING SCHEDULE
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)		
18. SUPPLEMENTARY NOTES		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) GOODNESS OF FIT WEIBULL DISTRIBUTION EXTREME VALUE DISTRIBUTION "EMPIRICAL" STOCHASTIC PROCESS WEAK CONVERGENCE TEST STATISTICS		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) In this note we consider some test statistics based on the sample distribution function for testing the null hypothesis that a random sample belongs to a Weibull distribution with unknown scale and shape parameters. A foundation for testing such a hypothesis is provided by the fact that the logarithm of a Weibull random variable has an extreme value distribution with a location and a scale parameter, and by some recent results of Durbin (1973) and of Serfling and Wood (1976). These results pertain to		

DD FORM 1473  
1 JAN 73EDITION OF 1 NOV 65 IS OBSOLETE  
S/N 0102-014-6601NONE  
SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

NONE

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

20. ABSTRACT (cont'd)

the weak convergence of an associated "empirical" stochastic process, under the null hypothesis. The asymptotic distribution of the empirical process serves as a basis for Monte Carlo studies for determining the appropriate critical points of the test statistics. We shall give some results on comparing the power of our tests and a test due to Mann, Scheuer, and Fertig (1973).

ACCESSION for	
NTIS	Write Section <input checked="" type="checkbox"/>
DDC	B.H. Section <input type="checkbox"/>
UNANNOUNCED	<input type="checkbox"/>
JUL 1 1973	
BY	
DISTRIBUTION AVAILABILITY CODES	
0	SPECIAL
A	

NONE

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)



THE GEORGE WASHINGTON UNIVERSITY  
School of Engineering and Applied Science  
Institute for Management Science and Engineering

Abstract  
of  
Serial T-363  
14 October 1977

ON SOME GOODNESS OF FIT TESTS FOR THE WEIBULL  
DISTRIBUTION WITH ESTIMATED PARAMETERS

by

Mahesh Chandra  
Nozer D. Singpurwalla

In this note we consider some test statistics based on the sample distribution function for testing the null hypothesis that a random sample belongs to a Weibull distribution with unknown scale and shape parameters. A foundation for testing such a hypothesis is provided by the fact that the logarithm of a Weibull random variable has an extreme value distribution with a location and a scale parameter, and by some recent results of Durbin (1973) and of Serfling and Wood (1976). These results pertain to the weak convergence of an associated "empirical" stochastic process, under the null hypothesis. The asymptotic distribution of the empirical process serves as a basis for Monte Carlo studies for determining the appropriate critical points of the test statistics. We shall give some results on comparing the power of our tests and a test due to Mann, Scheuer, and Fertig (1973).

THE GEORGE WASHINGTON UNIVERSITY  
School of Engineering and Applied Science  
Institute for Management Science and Engineering

ON SOME GOODNESS OF FIT TESTS FOR THE WEIBULL  
DISTRIBUTION WITH ESTIMATED PARAMETERS

by

Mahesh Chandra  
Nozer D. Singpurwalla

1. Introduction

The two-parameter Weibull distribution has found many applications in the engineering and in the biological sciences. For instance, it has been used by Cook, Doll and Fellingham (1969) and by Doll (1971), to describe the observed age distribution of many human cancers. Its use for describing failures of electrical and mechanical components is well documented in the reliability literature.

In this note we address ourselves to a fundamental problem involving any application of the Weibull distribution. We wish to test the null hypothesis that a given random sample belongs to a Weibull distribution with unknown parameters. Of the several methods for testing "goodness of fit," those based on the sample distribution function happen to be the most popular. We shall present tables of critical values for testing the null hypothesis in question, and also give some results comparing the power of our tests and a test due to Mann, Scheuer and Fertig (1973).

A foundation for developing our tables of critical values is the recently given theory by Durbin (1973), and by Serfling and Wood (1976) on the weak convergence of an "empirical" stochastic process. This



stochastic process is based on the sample distribution function and estimates of the unknown parameters. The statistics that we discuss can be represented as well-behaved functionals of this empirical process. Thus, the asymptotic distributions of the relevant test statistics can be obtained as the distributions of the corresponding functionals of the limiting process. The above ideas will be made clear in the following text.

## 2. Preliminaries

The two-parameter Weibull distribution is given by

$$F(t) = 1 - \exp\left[-\left(\frac{t}{\delta}\right)^\beta\right], \quad t \geq 0$$

$$= 0, \quad \text{otherwise;} \quad (2.1)$$

the scale parameter  $\delta$  and the shape parameter  $\beta$  are both assumed to be positive.

If we make the transformation  $X = -\ln T$ , where  $T$  has a two-parameter Weibull distribution, then the distribution of  $X$  is called the extreme value distribution. It is given by

$$F(x) = \exp\left(-\exp - \left(\frac{x-a}{b}\right)\right), \quad (2.2)$$

where  $a = -\ln \delta$  and  $b = \frac{1}{\beta}$ . We note that  $a$  and  $b$  are, respectively, the location and the scale parameters of the extreme value distribution.

The tests that we discuss in this paper are based on the extreme value distribution. To make a test of fit to the Weibull distribution we shall first take the negative of the natural logarithms of the supposed Weibull data. Thus, we wish to consider the case of testing whether the distribution of a random sample  $X_1, X_2, \dots, X_n$  is an extreme value distribution with unknown location parameter  $a$  and unknown scale parameter  $b$ . Specifically, we wish to test the "null hypothesis"

$$H_0 : F(x) = \exp\left[-\exp - \left(\frac{x-a}{b}\right)\right],$$

for all  $x$  and for some  $(a,b)$ .

When  $a$  and  $b$  are specified, then  $H_0$  is "simple," and our test reduces to testing the hypothesis that the independent random variables

$$G\left[\frac{X_i - a}{b}\right] = \exp\left[-\exp - \left(\frac{X_i - a}{b}\right)\right], \quad 1 \leq i \leq n,$$

have a common uniform  $(0,1)$  distribution. The Kolmogorov-Smirnov test is based on the statistic

$$n^{1/2} \sup_{0 \leq t \leq 1} |G_n(t) - t|, \quad (2.3)$$

where

$$G_n(t) = \frac{1}{n} \sum_{i=1}^n I\left(G\left[\frac{X_i - a}{b}\right] \leq t\right), \quad 0 \leq t \leq 1, \quad (2.4)$$

where  $I(E)$  denotes the indicator of the event  $E$ . Under the null hypothesis, the "empirical" stochastic process

$$W_n(t) = n^{1/2} [G_n(t) - t], \quad 0 \leq t \leq 1 \quad (2.5)$$

satisfies

$$W_n \xrightarrow{d} W^0 \quad \text{in } \mathcal{D}[0,1], \quad (2.6)$$

where  $\xrightarrow{d}$  denotes convergence in distribution and  $W^0$  denotes the Gaussian process determined by

$$E[W^0(t)] = 0, \quad 0 \leq t \leq 1$$

and

$$E[W^0(s)W^0(t)] = \min(s,t) - st, \quad 0 \leq s, t \leq 1.$$

$\mathcal{D}[0,1]$  denotes the space of functions on  $[0,1]$  which are right-continuous and have left-hand limits.

In the following section we present some results on an analogous test statistic for the case  $H_0$  composite. These results will serve as a basis for developing the tables of critical values.

### 3. The Convergence Theorem and the Test Statistic

When  $a$  and  $b$  are not specified, that is, when  $H_0$  is composite, we consider an analogous approach based on  $(\hat{a}_n, \hat{b}_n)$ , the maximum likelihood estimators of  $(a, b)$ . We set

$$Y_{n,i} = \frac{X_i - \hat{a}_n}{\hat{b}_n}, \quad 1 \leq i \leq n,$$

and analogous to  $G_n$  and  $W_n$  we define

$$H_n(t) = \frac{1}{n} \sum_{i=1}^n I[G(Y_{n,i}) \leq t], \quad 0 \leq t \leq 1$$

and

$$V_n(t) = n^{1/2} [H_n(t) - t], \quad 0 \leq t \leq 1.$$

Our theorem pertains to the "empirical" stochastic process  $V_n(t)$ , and is analogous to the result given by Equation (2.6). However, before stating the convergence theorem, we will have to introduce the following notation given in Durbin (1973), and verify that his assumptions (conditions) are satisfied.

Let us denote by  $\theta$  the vector  $[a, b]'$ , and let  $\theta_0$  be any conveniently chosen value of  $\theta$ . We state below a verification of the required conditions.

Condition A: The distribution  $G(x, \theta_0)$  has a density  $f(x, \theta_0)$  such that, for almost all  $x$ , the vector  $\partial \log f(x, \theta_0) / \partial \theta_0$  exists, and satisfies

$$E \left( \frac{\partial \log f(x, \theta_0)}{\partial \theta_0}, \frac{\partial \log f(x, \theta_0)}{\partial \theta_0'} \right) = J,$$

where  $J$  is finite and positive definite.

Condition B: Let  $\hat{\theta}_n$  be the maximum likelihood estimator of  $\theta$ ; that is,  $\hat{\theta}_n = [\hat{a}_n, \hat{b}_n]'$ . Then, it is well known (cf. Cramer (1946)) that

$$n^{1/2} (\hat{\theta}_n - \theta_0) = \frac{1}{n^{1/2}} J^{-1} \sum_{i=1}^n \frac{\partial \log f(x_i, \theta_0)}{\partial \theta_0} + \varepsilon_n,$$

where  $\varepsilon_n \rightarrow 0$ , in probability.

Condition C: Let  $N$  be the closure of a neighborhood of  $\theta_0$ . Let  $g(t, \theta) = \partial G(x, \theta) / \partial \theta$  when this is expressed as a function of  $t$  by means of the transformation  $t = G(x, \theta)$ ; let  $g(t) = g(t, \theta_0)$ . The vector function  $g(t, \theta)$  is continuous in  $(\theta, t)$  for all  $\theta \in N$ , and  $0 \leq t \leq 1$ .

Theorem 3.1: By virtue of Conditions A, B, and C, the "empirical" process  $V_n$  determined by the extreme value distribution  $G\left[\frac{x-a_n}{b_n}\right]$ , with  $(a_n, b_n)$  the maximum likelihood estimators, is such that

$$V_n \xrightarrow{d} V^0 \quad \text{in } \mathcal{D}[0,1],$$

where  $V^0$  is a Gaussian process determined by

$$E[V^0(t)] = 0, \quad 0 \leq t \leq 1$$

and

$$E[V^0(s)V^0(t)] = \min(s, t) - st - g(s)' J^{-1} g(t), \quad 0 \leq s, t \leq 1. \quad (3.1)$$

Proof: Follows from Durbin (1973). //

If we choose  $\theta_0 = [0, 1]'$ , then it can be verified that  $g(t) = [t \log t, -t \log t \log(-\log t)]$ , and that

$$J^{-1} = \begin{vmatrix} 1.10867 & 0.257 \\ 0.257 & 0.60793 \end{vmatrix},$$

[cf. Johnson and Kotz (1970), p. 282]. Substituting the above into (3.1) we have the covariance of our Gaussian process

$$\begin{aligned} E[V^0(s)V^0(t)] = & \min(s,t) - st - 1.108(s \log s)(t \log t) \\ & + 0.257(s \log s)(t \log t \log(-\log t)) \\ & + 0.257(s \log s \log(-\log s)(t \log t)) \\ & - 0.60793(s \log s \log(-\log s) t \log t \log(-\log t)) , \quad 0 \leq s, t \leq 1 . \end{aligned} \quad (3.2)$$

The statistics of interest in connection with  $H_0$  are:

(i) the one-sided Kolmogorov statistic

$$D_n^+ = \sup_{0 \leq t \leq 1} V_n(t) , \quad (3.3)$$

$$D_n^- = -\inf_{0 \leq t \leq 1} V_n(t) , \quad (3.4)$$

(ii) the Kolmogorov-Smirnov statistic

$$D_n = \max(D_n^+, D_n^-) , \quad (3.5)$$

(iii) the Kupier statistic

$$V_n = D_n^+ + D_n^- , \quad (3.6)$$

(iv) the Cramer-Von Mises statistic

$$W_n^2 = \int_0^1 V_n^2(t) dt , \quad (3.7)$$

(v) the Watson statistic

$$U_n^2 = \int_0^1 V_n^2(t) dt - \left[ \int_0^1 V_n(t) dt \right]^2 , \quad (3.8)$$

and

(vi) the Anderson-Darling statistic

$$A_n^2 = \int_0^1 \frac{V_n^2(t)}{t(1-t)} dt . \quad (3.9)$$



Using the fact that if  $h(V_n(t))$  is a function of  $V_n(t)$  that is continuous with respect to the Skorokhod metric on  $D[0,1]$ , the limit laws of  $D_n^+$ ,  $D_n^-$ ,  $D_n$ ,  $V_n$ ,  $W_n^2$ ,  $U_n^2$ , and  $A_n^2$  under  $H_0$  are given, respectively, by the laws of the random variables

$$D^+ = \sup_{0 \leq t \leq 1} V^0(t), \quad (3.10)$$

$$D^- = -\inf_{0 \leq t \leq 1} V^0(t), \quad (3.11)$$

$$D = \max(D^+, D^-), \quad (3.12)$$

$$V = D^+ + D^-, \quad (3.13)$$

$$W^2 = \int_0^1 (V^0(t))^2 dt, \quad (3.14)$$

$$U^2 = \int_0^1 (V^0(t) dt)^2 - \left[ \int_0^1 V^0(t) dt \right]^2, \quad (3.15)$$

and

$$A^2 = \lim_{\epsilon \rightarrow 0} \int_{0+\epsilon}^{1-\epsilon} \frac{(V^0(t))^2}{t(1-t)} dt. \quad (3.16)$$

The above results follow as a consequence of the continuous mapping theorem of Billingsley (1968). They provide a basis for Monte Carlo studies of the null hypothesis asymptotic distributions of the statistics discussed above.

#### 4. Sampling Distributions of the Approximate Test Statistics

Monte Carlo methods were used to simulate the distribution of the limiting random variables given in Equations (3.10) through (3.16). Following Serfling and Wood (1976) we approximate the Gaussian process  $V^0$  by its finite-dimensional distributions, corresponding to an evaluation of the process at 29, 99, and 119 equally-spaced points in the unit

interval. One thousand multivariate normal random vectors with the covariance given by Equation (3.2) were generated using a program from the International Mathematical and Statistical Library. The empirical distributions of the supremum, the infimum, and the difference between the supremum and the infimum of the resulting multivariate normal vectors were then tabulated, thus approximating the limit laws of  $D_n^+$ ,  $D_n^-$ ,  $D_n$ , and  $V_n$ . Since the differences in the observed quantiles corresponding to the finite-dimensional distributions of  $V^0$  at 29, 99, and 119 equally-spaced points diminished, the approximating procedure was terminated at 119 equally-spaced points. The asymptotic distributions of  $W^2$ ,  $U^2$ , and  $A^2$  were obtained by using numerical integration techniques. For this we used Subroutine QSF from the IBM Scientific Subroutine Package. The various sample quantiles for the generated frequency distributions are shown in Table 4.1.

#### 5. The Mann-Scheuer-Fertig (MSF) Test

The only other known procedure for testing goodness of fit for the Weibull that is not based on the empirical distribution function is a test proposed by Mann, Scheuer, and Fertig (1973).

The MSF test is based on a statistic  $S$ , and can be used for censored as well as uncensored samples. However, the percentage points of  $S$  and certain quantities that are used in calculating  $S$  are available only for sample sizes of up to 25. However, along with a modification given by Stephens (1977), the test statistics we discuss can be used for any sample size.

For a sample of size  $n$ , censored at  $m$ , the statistic  $S$  is defined as

$$S = \frac{\sum_{i=[m/2]+1}^{m-1} (X_{i+1} - X_i) / [E(Y_{i+1}) - E(Y_i)]}{\sum_{i=1}^{m-1} (X_{i+1} - X_i) / [E(Y_{i+1}) - E(Y_i)]},$$

where  $Y_i = \frac{x_i - a}{b}$  and  $[r]$  denotes the greatest integer contained in  $r$ . Mann, Scheuer, and Fertig give percentage points of  $S$  and the values of the quantities  $[E(Y_{i+1}) - E(Y_i)]$  for samples of size 3 to 25.

## 6. Power Comparisons

In order to evaluate the effectiveness of the tests discussed before, we evaluate their power, against the lognormal distribution as an alternative. The lognormal distribution is chosen because it appears to be a natural competitor to a Weibull distribution. The power comparisons were made numerically. For this random samples of size 20, 25, and 30, respectively, were generated from a lognormal (normal) distribution with parameters -0.5 (mean) and 1.00 (variance), respectively.

Maximum likelihood estimators of the parameters  $a$  and  $b$  of the extreme value distribution were obtained by numerically solving the following equations simultaneously:

$$\hat{b} = \sum_j X_j / n - \left[ \sum_j X_j \exp(-X_j / \hat{b}) \right] \left[ \sum_j \exp(-X_j / \hat{b}) \right]^{-1} \quad (6.1)$$

and

$$\hat{a} = -\hat{b} \log \left[ \sum_j \exp(-X_j / \hat{b}) / n \right]. \quad (6.2)$$

The results of our power comparisons are shown in Tables 6.1, 6.2, and 6.3, and these are based on 1000 replicates. Based on this limited experiment, it appears that for samples of sizes 20 and 25, the MSF test has better power. For samples of size 30, the MSF test could not be used, and the Anderson-Darling test appears to have better power.

## 7. Concluding Remarks

After finishing the work on this report we were informed that Stephens (1977) has also obtained asymptotic percentage points for the statistics  $W^2$ ,  $U^2$ , and  $A^2$ . Stephens also gives a necessary modification so as to use these statistics for a finite sample sizes. Even though

TABLE 6.1  
POWER COMPARISON: WEIBULL vs. LOGNORMAL  
SAMPLE SIZE  $n = 20$

Level of Significance	Kolmogorov-Smirnov			Kupier $V$	Cramer von Mises $W^2$	Watson $U^2$	Anderson Darling $A^2$	Mann Scheuer Fertig $S$
	$D^+$	$D^-$	$D$					
0.01	0.008	0.101	0.075	0.102	0.080	0.082	0.100	0.105
0.025	0.023	0.167	0.114	0.131	0.144	0.140	0.177	Not Available
0.05	0.046	0.236	0.171	0.209	0.219	0.211	0.238	0.265
0.10	0.084	0.418	0.249	0.303	0.322	0.321	0.354	0.432

TABLE 6.2  
POWER COMPARISON: WEIBULL vs. LOGNORMAL  
SAMPLE SIZE  $n = 25$

Level of Significance	Kolmogorov-Smirnov			Kupier V	Cramer von Mises $W^2$	Watson $U^2$	Anderson Darling $A^2$	Mann Scheuer Fertig S
	$D^+$	$D^-$	D					
0.01	0.019	0.145	0.121	0.141	0.131	0.132	0.166	0.153
0.025	0.034	0.196	0.155	0.189	0.196	0.190	0.240	Not Available
0.05	0.072	0.263	0.206	0.255	0.279	0.264	0.310	0.389
0.10	0.116	0.457	0.286	0.360	0.364	0.365	0.411	0.533



TABLE 6.3  
POWER COMPARISON: WEIBULL vs. LOGNORMAL  
SAMPLE SIZE  $n = 30$

Level of Significance	Kolmogorov-Smirnov			Kupier V	Cramer von Mises $w^2$	Watson $u^2$	Anderson Darling $A^2$	Mann Scheuer Fertig S
	$D^+$	$D^-$	D					
0.01	0.021	0.173	0.137	0.161	0.151	0.145	0.183	Not Available
0.025	0.042	0.252	0.185	0.210	0.231	0.219	0.286	Not Available
0.05	0.082	0.331	0.256	0.296	0.333	0.319	0.369	Not Available
0.10	0.132	0.521	0.355	0.412	0.421	0.418	0.475	Not Available

our approach is different, it is encouraging to note that our results seem to be in good agreement with those of Stephens. A comparison of the asymptotic points we obtained with those of Stephens is given in Table 7.1. Stephens has made no power comparisons, and since our results agree quite well with his, we conclude that our power comparisons remain valid.

#### ACKNOWLEDGMENT

We are grateful to Professor Robert J. Serfling of the Florida State University, who was kind enough to provide us with a preprint of his paper. Our work is motivated by Section 4 of his paper.

TABLE 7.1  
COMPARISON OF UPPER TAIL PERCENTAGE POINTS OF  $W^2$ ,  $U^2$ , AND  $A^2$  STATISTICS

Statistic	$\alpha = 0.75$	$\alpha = 0.90$	$\alpha = 0.95$	$\alpha = 0.975$	$\alpha = 0.99$
	Table 4.1   Stephens	Table 4.1   Stephens	Table 4.1   Stephens	Table 4.1   Stephens	Table 4.1   Stephens
$W^2$	0.073   0.073	0.105   0.102	0.123   0.124	0.147   0.146	0.175   0.175
$U^2$	0.069   0.070	0.098   0.097	0.117   0.117	0.140   0.138	0.164   0.165
$A^2$	0.457   0.474	0.623   0.637	0.746   0.757	0.849   0.877	0.991   1.038

## REFERENCES

- BILLINGSLEY, PATRICK (1968). Convergence of Probability Measures.  
Wiley, New York.
- COOK, P., R. DOLL and S. A. FELLINGHAM (1969). A mathematical model  
for the age distribution of cancer in man. Internat. J. Cancer  
 $\sqrt[4]{}$  93-112.
- CRAMER, H. (1946). Mathematical Methods of Statistics. Princeton  
University Press, Princeton, New Jersey.
- DOLL, R. (1971). The age distribution of cancer: implications for models  
of carcinogens. J. Roy. Statist. Soc. Ser. A  $\sqrt[134]{}$  133-166.
- DURBIN, J. (1973). Weak convergence of the sample distribution function  
when parameters are estimated. Ann. Statist.  $\sqrt[1]{}$  279-290.
- JOHNSON, N. L. and S. KOTZ (1970). Continuous Univariate Distributions,  
 $\sqrt[1]{}$ , Houghton-Mifflin Company, Boston.
- MANN, NANCY R., E. M. SCHEUER and K. W. FERTIG (1973). A new goodness  
of fit test for the two parameter Weibull or extreme value distri-  
bution with unknown parameters. Comm. in Statistics  $\sqrt[2]{}$  (5) 383-400.
- SERFLING, R. J. and C. L. WOOD (1976). On null hypothesis limiting dis-  
tributions of Kolmogorov-Smirnov type statistics with estimated  
location and scale parameters. Technical report, Florida State  
University, Tallahassee, Florida.
- STEPHENS, M. A. (1977). Goodness-of-fit for the extreme value distribution.  
Technical Report No. 247, Department of Statistics, Stanford University,  
Stanford, California.

THE GEORGE WASHINGTON UNIVERSITY  
Institute for Management Science and Engineering  
Distribution List for Technical Papers

The George Washington University  
Office of Sponsored Research  
Library  
Vice President H. F. Bright  
Dean Harold Liebowitz  
Mr. J. Frank Doubleday

ONR  
Chief of Naval Research  
(Codes 200, 430D, 1021P)  
Resident Representative

OPNAV  
OP-40  
DCNO, Logistics  
Navy Dept Library  
OP-911  
OP-964

Naval Aviation Integrated Log Support

NAVCOSSACT

Naval Cmd Sys Sup Activity Tech Library

Naval Electronics Lab Library

Naval Facilities Eng Cmd Tech Library

Naval Ordnance Station  
Louisville, Ky.  
Indian Head, Md.

Naval Ordnance Sys Cmd Library

Naval Research Branch Office  
Boston  
Chicago  
New York  
Pasadena  
San Francisco

Naval Research Lab  
Tech Info Div  
Library, Code 2029 (ONRL)

Naval Ship Engng Center  
Philadelphia, Pa.  
Hyattsville, Md.

Naval Ship Res & Dev Center

Naval Sea Systems Command  
Tech Library  
Code 073

Naval Supply Systems Command  
Library  
Capt W. T. Nash

Naval War College Library  
Newport

BUPERS Tech Library

FMSO

Integrated Sea Lift Study

USN Ammo Depot Earle

USN Postgrad School Monterey  
Library  
Dr. Jack R. Borsting  
Prof C. R. Jones

US Marine Corps  
Commandant  
Deputy Chief of Staff, R&D

Marine Corps School Quantico  
Landing Force Dev Ctr  
Logistics Officer

Armed Forces Industrial College

Armed Forces Staff College

Army War College Library  
Carlisle Barracks

Army Cmd & Gen Staff College

US Army HQ  
LTC George L. Slyman  
Army Trans Mat Command

Army Logistics Mgmt Center  
Fort Lee

Commanding Officer, USALDSRA  
New Cumberland Army Depot

US Army Inventory Res Ofc  
Philadelphia

HQ, US Air Force  
AFADS-3

Griffiss Air Force Base  
Reliability Analysis Center

Maxwell Air Force Base Library

Wright-Patterson Air Force Base  
HQ, AF Log Command  
Research Sch Log

Defense Documentation Center

National Academy of Science  
Maritime Transportation Res Board Library

National Bureau of Standards  
Dr E. W. Cannon  
Dr Joan Rosenblatt

National Science Foundation

National Security Agency

WSEG

British Navy Staff

Logistics, OR Analysis Establishment  
National Defense Hdqtrs, Ottawa

American Power Jet Co  
George Chernowitz

ARCON Corp

General Dynamics, Pomona

General Research Corp  
Dr Hugh Cole  
Library

Planning Research Corp  
Los Angeles

Rand Corporation  
Library

Carnegie-Mellon University  
Dean H. A. Simon  
Prof G. Thompson

Case Western Reserve University  
Prof B. V. Dean  
Prof John R. Isbell  
Prof M. Mesarovic  
Prof S. Zacks

Cornell University  
Prof R. E. Bechhofer  
Prof R. W. Conway  
Prof J. Kiefer  
Prof Andrew Schultz, Jr.

Cowles Foundation for Research  
Library  
Prof Herbert Scarf  
Prof Martin Shubik

Florida State University  
Prof R. A. Bradley

Harvard University  
Prof K. J. Arrow  
Prof W. G. Cochran  
Prof Arthur Schleifer, Jr.

New York University  
Prof O. Morgenstern

Princeton University  
Prof A. W. Tucker  
Prof J. W. Tukey  
Prof Geoffrey S. Watson



**Purdue University**  
 Prof S. S. Gupta  
 Prof H. Rubin  
 Prof Andrew Whinston

**Stanford**  
 Prof T. W. Anderson  
 Prof G. B. Dantzig  
 Prof F. S. Hillier  
 Prof D. L. Iglehart  
 Prof Samuel Karlin  
 Prof G. J. Lieberman  
 Prof Herbert Solomon  
 Prof A. F. Veinott, Jr.

**University of California, Berkeley**  
 Prof R. E. Barlow  
 Prof D. Gale  
 Prof Rosedith Sitgreaves  
 Prof L. M. Tichvinsky

**University of California, Los Angeles**  
 Prof J. R. Jackson  
 Prof Jacob Marschak  
 Prof R. R. O'Neill  
 Numerical Analysis Res Librarian

**University of North Carolina**  
 Prof W. L. Smith  
 Prof M. R. Leadbetter

**University of Pennsylvania**  
 Prof Russell Ackoff  
 Prof Thomas L. Saaty

**University of Texas**  
 Prof A. Charnes

**Yale University**  
 Prof F. J. Anscombe  
 Prof I. R. Savage  
 Prof M. J. Sobel  
 Dept of Admin Sciences

Prof Z. W. Birnbaum  
 University of Washington

Prof B. H. Bissinger  
 The Pennsylvania State University

Prof Seth Bonder  
 University of Michigan

Prof G. E. P. Box  
 University of Wisconsin

Dr. Jerome Bracken  
 Institute for Defense Analyses

Prof H. Chernoff  
 MIT

Prof Arthur Cohen  
 Rutgers - The State University

Mr Wallace M. Cohen  
 US General Accounting Office

Prof C. Derman  
 Columbia University

Prof Paul S. Dwyer  
 Mackinaw City, Michigan

Prof Saul I. Gass  
 University of Maryland

Dr Donald P. Gaver  
 Carmel, California

Dr Murray A. Geisler  
 Logistics Mgmt Institute

Prof J. F. Hannan  
 Michigan State University

Prof H. O. Hartley  
 Texas A & M Foundation

Mr Gerald F. Hein  
 NASA, Lewis Research Center

Prof W. M. Hirsch  
 Courant Institute

Dr Alan J. Hoffman  
 IBM, Yorktown Heights

Dr Rudolf Husser  
 University of Bern, Switzerland

Prof J. H. K. Kao  
 Polytech Institute of New York

Prof W. Kruskal  
 University of Chicago

Prof C. E. Lemke  
 Rensselaer Polytech Institute

Prof Loynes  
 University of Sheffield, England

Prof Steven Nahmias  
 University of Pittsburgh

Prof D. B. Owen  
 Southern Methodist University

Prof E. Parzen  
 State University New York, Buffalo

Prof H. O. Posten  
 University of Connecticut

Prof R. Remage, Jr.  
 University of Delaware

Dr Fred Rigby  
 Texas Tech College

Mr David Rosenblatt  
 Washington, D. C.

Prof M. Rosenblatt  
 University of California, San Diego

Prof Alan J. Rowe  
 University of Southern California

Prof A. H. Rubenstein  
 Northwestern University

Dr M. E. Salvesson  
 West Los Angeles

Prof Edward A. Silver  
 University of Waterloo, Canada

Prof R. M. Thrall  
 Rice University

Dr S. Vajda  
 University of Sussex, England

Prof T. M. Whitin  
 Wesleyan University

Prof Jacob Wolfowitz  
 University of Illinois

Mr Marshall K. Wood  
 National Planning Association

Prof Max A. Woodbury  
 Duke University

# THE GEORGE WASHINGTON UNIVERSITY

BENEATH THIS PLAQUE  
IS BURIED

A VAULT FOR THE FUTURE  
IN THE YEAR 2056

THE STORY OF ENGINEERING IN THIS YEAR OF THE PLACING OF THE VAULT AND  
ENGINEERING HOPES FOR THE TOMORROWS AS WRITTEN IN THE RECORDS OF THE  
FOLLOWING GOVERNMENTAL AND PROFESSIONAL ENGINEERING ORGANIZATIONS AND  
THOSE OF THIS GEORGE WASHINGTON UNIVERSITY.

BOARD OF COMMISSIONERS, DISTRICT OF COLUMBIA  
UNITED STATES ATOMIC ENERGY COMMISSION  
DEPARTMENT OF THE ARMY, UNITED STATES OF AMERICA  
DEPARTMENT OF THE NAVY, UNITED STATES OF AMERICA  
DEPARTMENT OF THE AIR FORCE, UNITED STATES OF AMERICA  
NATIONAL ADVISORY COMMITTEE FOR AERONAUTICS  
NATIONAL BUREAU OF STANDARDS, U.S. DEPARTMENT OF COMMERCE  
AMERICAN SOCIETY OF CIVIL ENGINEERS  
AMERICAN INSTITUTE OF ELECTRICAL ENGINEERS  
THE AMERICAN SOCIETY OF MECHANICAL ENGINEERS  
THE SOCIETY OF AMERICAN MILITARY ENGINEERS  
AMERICAN INSTITUTE OF MINING & METALLURGICAL ENGINEERS  
DISTRICT OF COLUMBIA SOCIETY OF PROFESSIONAL ENGINEERS, INC.  
THE INSTITUTE OF RADIO ENGINEERS, INC.  
THE CHEMICAL ENGINEERS CLUB OF WASHINGTON  
WASHINGTON SOCIETY OF ENGINEERS  
FAULKNER, KINGSBURY & STENHOUSE - ARCHITECTS  
CHARLES H. TOMPKINS COMPANY - BUILDERS  
SOCIETY OF WOMEN ENGINEERS  
NATIONAL ACADEMY OF SCIENCES, NATIONAL RESEARCH COUNCIL

THE PURPOSE OF THIS VAULT IS INSPIRED BY AND IS DEDICATED TO  
**CHARLES HOOK TOMPKINS**, DOCTOR OF ENGINEERING  
BECAUSE OF HIS ENGINEERING CONTRIBUTIONS TO THIS UNIVERSITY, TO HIS  
COMMUNITY, TO HIS NATION, AND TO OTHER NATIONS.

BY THE GEORGE WASHINGTON UNIVERSITY.

ROBERT V. FLEMING  
CHAIRMAN OF THE BOARD OF TRUSTEES

CLOYD H. MARVIN  
PRESIDENT

JUNE THE TWENTIETH  
1956

To cope with the expanding technology, our society must be assured of a continuing supply of rigorously trained and educated engineers. The School of Engineering and Applied Science is completely committed to this objective.